

Soft and collinear parton radiation in heavy quark production*

P. M. Nadolsky¹, N. Kidonakis^{1,2}, F. Olness¹, C.-P. Yuan³

¹*Department of Physics, Southern Methodist University, Dallas, TX 75275-0175*

²*Department of Physics and Astronomy, University of Rochester, Rochester, NY 14627-0171*

³*Department of Physics & Astronomy, Michigan State University, East Lansing, MI 48824*

When the energy of the heavy quark is comparable with its mass, it is natural to attribute this heavy quark to the hard part of the reaction. At large energies, this approach is impractical due to large logarithms from intensive QCD radiation affecting both inclusive and differential observables. We present a formalism for all-order summation of such logarithms and reliable description of heavy-quark distributions at all energies. As an illustration, we calculate angular distributions of B mesons produced in neutral-current events at large momentum transfers at the ep collider HERA.

In recent years, significant attention was dedicated to exploring properties of heavy-flavor hadrons produced in lepton-nucleon deep inelastic scattering (DIS). On the experimental side, the Hadron-Electron Ring Accelerator at DESY (HERA) has generated a large amount of data on the production of charm and bottom mesons in neutral current interactions (cf. [1] and references therein). At present energies (about 300 GeV in the ep c.m. frame), a substantial charm production cross section is observed in a wide range of Bjorken x and photon virtualities Q^2 , and charm quarks contribute up to 30% to the DIS structure functions.

On the theory side, Perturbative Quantum Chromodynamics (PQCD) provides a natural framework for the description of the heavy-flavor production. Due to large masses M of charm and bottom quarks ($M^2 \gg \Lambda_{QCD}^2$), the renormalization scale can be always chosen in a region where the effective strong coupling α_S is small. Despite the smallness of α_S , perturbative calculations in the presence of heavy flavors are not without intricacies. In particular, care in the choice of a proper factorization scheme is essential for the efficient separation of short- and long-distance contributions to the heavy quark cross section. Recent reviews of the theoretical status can be found in Refs. [2, 3, 4]. The key issue here is whether, for a given renormalization and factorization scale $\mu \sim Q$, the heavy quarks of the N -th flavor are treated as *partons* in the incoming proton, *i.e.*, whether one calculates the QCD beta-function using N active flavors and introduces a parton distribution function (PDF) for the N -th flavor. A related, but separate, issue is whether the mass of the heavy quark can be neglected in the hard cross section and/or parton distribution function without ruining the accuracy of the calculation. Currently, several factorization schemes [5, 6, 7, 8, 9, 10, 11, 12] are available that differ in the treatment of these issues; of those, the variable flavor number schemes reduce to the conventional massless $\overline{MS}$ factorization scheme in the limit $Q^2 \gg M^2$. The general proof of the factorization in the presence of nonzero heavy quark masses was given in Ref. [13].

In the near future, we expect the quality of the heavy flavor data to greatly improve. By 2006, the upgraded ep collider HERA will accumulate an integrated luminosity of 1 fb^{-1} , *i.e.*, more than eight times the final integrated luminosity from its previous runs. Studies of heavy quarks in ep collisions are also envisioned at the proposed Electron Ion Collider [14] and THERA [15]. By the end of these experiments, a large amount of information will be available both for the inclusive observables (like the charm component of the DIS structure functions) and differential distributions.

In our work [16], we address a theoretical formalism for the description of heavy-quark differential distributions both near the heavy-quark mass threshold ($Q^2 \approx M^2$) and far beyond it ($Q^2 \gg M^2$). Our attention is primarily focused on the angular or transverse momentum distributions of the final-state heavy mesons. The behavior of such distributions is quite complex, since three distinct momentum scales (the heavy quark mass M , the photon virtuality Q , and the transverse momentum of the heavy quark) are involved. As a

*Talk presented at the American Physical Society's 2002 Meeting of The Division of Particles and Fields (Williamsburg, VA, 2002)




Figure 1: The neutral current semi-inclusive production $e + p \rightarrow e + H + X$ of heavy mesons in the $\gamma^* p$ c.m. reference frame. q^μ , p^μ , and p_H^μ are momenta of the virtual photon, proton, and the heavy meson, respectively. The polar angle θ_H of the heavy meson defines the scale q_T^2 discussed in the text. The resummation effects considered here are important in the limit $\theta_H \rightarrow 0$, *i.e.*, when the final-state heavy meson closely follows the direction of its escape in the lowest-order flavor-excitation process $\gamma^* + q \rightarrow q$.

result, large logarithms of ratios of two very different scales tend to appear in the fixed-order perturbative cross sections in some regions of the phase space. These logarithms break the convergence of the series in the strong coupling α_S . To obtain stable theoretical predictions, they have to be summed through all orders of α_S . The “resummation” is realized by the reorganization of factorized cross sections and solution of equations for the gauge and renormalization group invariance.

The inclusive and semi-inclusive heavy-flavor cross sections may contain large logarithms of two kinds. The logarithms $\ln(Q^2/M^2)$ appear already in the inclusive observables and are resummed in the heavy-quark PDFs. The resummation of this type constitutes the essence of the massive variable flavor number factorization schemes, as, for instance, the ACOT scheme [6]. Logarithms of the form $(\alpha_S^n/q_T^2) \ln^m(q_T^2/Q^2)$, $0 \leq m \leq 2n - 1$ appear at $Q^2 \gg M^2, q_T^2$ when calculating angular differential distributions. Here $q_T \equiv p_T/z$ is the transverse momentum p_T of the heavy meson H rescaled by the final-state fragmentation variable $z \equiv (p \cdot p_H)/(p \cdot q)$; p^μ , q^μ , and p_H^μ are the momenta of the initial-state proton, virtual photon, and heavy meson, respectively. The limit $q_T^2 \ll Q^2$ is equivalent to the limit $\theta_H \rightarrow 0$, where θ_H is the polar angle of H in the $\gamma^* p$ c.m. reference frame (cf. Fig. 1). If M is neglected ($M^2 \ll Q^2$), q_T is simply related to θ_H :

$$q_T^2 = Q^2 \left(\frac{1}{x} - 1 \right) \frac{1 - \cos \theta_H}{1 + \cos \theta_H} \longrightarrow Q^2 \left(\frac{1}{x} - 1 \right) \left(\frac{\theta_H^2}{4} + \dots \right) \text{ as } \theta_H \rightarrow 0.$$

The logarithms $\ln(q_T^2/Q^2)$ originate from the incomplete cancellation of soft singularities and factorization of quasi-collinear singularities in the fixed-order cross sections.¹ The formalism for their summation was proposed by Collins, Soper, and Sterman (CSS) to describe angular correlations in e^+e^- hadroproduction [18, 19] and transverse momentum distributions in the Drell-Yan process [20]. Recently this formalism was extended to describe semi-inclusive DIS (SIDIS) production of practically *massless* hadrons [21, 22, 23]. The resummation in massless SIDIS serves as a high-energy limit for our calculation.

¹ A discussion of integrable singularities in the fixed-order differential distributions for the semi-inclusive DIS charm production can be found in Ref. [17].




Figure 2: Balance of various terms in the ACOT scheme and resummed cross section. Graphs show q_T on the x -axis and $d\sigma/dq_T^2$ on the y -axis.

The key result of the CSS formalism is that all logarithms due to the soft radiation in the high-energy limit can be summed into a Sudakov exponent. This result can be summarized in the following master equation:

$$\frac{d\sigma}{dQ^2 dq_T^2} = \frac{\sigma_0}{s} \int \frac{d^2b}{(2\pi)^2} e^{i\vec{q}_T \cdot \vec{b}} (C^{in} \otimes f) (C^{out} \otimes d) e^{-S} + \sigma_{FO} - \sigma_{ASY}. \quad (1)$$

Here b is the impact parameter conjugate to q_T , f and d are parton distributions and fragmentation functions, respectively. C^{in} , C^{out} contain perturbative corrections to contributions from the incoming and outgoing hadronic jets, respectively. The factor e^{-S} is the Sudakov exponential, which includes an all-order sum of perturbative logarithms $\ln^m(q_T^2/Q^2)$ at $b \lesssim 1 \text{ GeV}^{-1}$ and nonperturbative contributions at $b \gtrsim 1 \text{ GeV}^{-1}$. Finally, σ_{FO} is the fixed-order expression for the considered cross section, while σ_{ASY} (*asymptotic piece*) is the perturbative expansion of the b -space integral up to the same order of α_S as in σ_{FO} . At small q_T , where terms $\ln^m(q_T^2/Q^2)$ are large, σ_{FO} cancels well with σ_{ASY} , so that the cross section (1) is approximated well by the b -space integral. At $q_T \gtrsim Q$, where the logarithms are no longer dominant, the b -space integral cancels with σ_{ASY} , so that the cross section (1) is equal to σ_{FO} up to higher-order corrections.

To extend the CSS resummation from the massless to the massive case, the logarithms of both types $\ln(Q^2/M^2)$ and $\ln(q_T^2/Q^2)$ should be resummed simultaneously. For the resummation of the logarithms $\ln(Q^2/M^2)$, we use the simplified ACOT factorization scheme [12, 13]. With regards to the logarithms $\ln(q_T^2/Q^2)$, we observe that the proof of the resummation formula in the original papers [18, 19, 20] does not rely on the dimensional regularization for the soft and collinear singularities appearing in individual Feynman diagrams. Although the dimensional regularization does provide a convenient way to regulate ultraviolet singularities in α_S and PDFs (and hence, defines the $\overline{MS}$ regularization and factorization schemes), it does not have to be invoked to regulate the singularities in the infrared region: other methods (*e.g.*, regularization with nonzero quark and gluon masses) will do just fine. As a result, the b -space integral is well-defined both for nonzero quark masses and in the limit $M/Q \rightarrow 0$; hence it cannot contain negative powers or logarithms of M with the exception of the collinear logarithms resummed in the PDFs or fragmentation functions. Also




Figure 3: The angular distributions of B mesons in the γ^*p c.m. frame close to the mass threshold ($Q = 5$ GeV, left figure) and at $Q^2 \gg M^2$ ($Q = 50$ GeV, right figure). In both cases, the cross sections are calculated for $x = 0.05$ and $M = 4.5$ GeV using the CTEQ5HQ PDFs [24] and Peterson fragmentation functions [25] with $\varepsilon = 0.0033$. The plots show the fixed-order cross section (long-dashed line), the b -space integral (dot-dashed line), the asymptotic piece (dotted line), and the full resummed cross section (solid line).

the quark masses regulate collinear singularities, so that at least some of the logarithms are not dominant in the threshold region. The remaining issue is about the accuracy of the approximation. The evolution equations used in the derivation of the b -space integral in Refs. [18, 19] did not keep track of the power-suppressed terms $\mathcal{O}((M/Q)^n)$, $n > 0$. As a result, the b -space integral does not necessarily approximate the small- q_T behavior of the cross section if M/Q is not negligible. However, the full Eq. (1) does a better job and approximates the cross section of the order $\mathcal{O}(\alpha_S^k)$ up to corrections of the order $\mathcal{O}(\alpha_S^{k+1}, M/Q)$. This happens because terms up to the order $\mathcal{O}(\alpha_S^k, M/Q)$ are canceled between the b -space integral and σ_{ASY} ; the terms $\ln^m(q_T^2/Q^2)$ that were not resummed (if any) cancel between σ_{FO} and σ_{ASY} .

Fig. 2 qualitatively illustrates the balance of various terms in the resummation formula in various regions of phase space for the lowest-order contributions. First, consider a fixed-order calculation in the simplified ACOT scheme near the threshold. In this region, the cross section $d\sigma/dq_T^2$ is well approximated by the $\mathcal{O}(\alpha_S)$ photon-gluon diagram (Fig. 2a). To this diagram, we add the lowest-order γ^*q term, which resums powers of $\ln(Q^2/M^2)$ and contributes at $q_T = 0$ (Fig. 2b). We also subtract the overlap between the two diagrams (Fig. 2c), which is approximately equal to, but not the same as, the γ^*q contribution. The resulting distribution (Fig. 2d) is close to the fixed-order result, but has discontinuities in the small q_T region. These discontinuities are amplified when Q increases.

In the q_T -resummed cross section, the fusion diagram still dominates near the threshold, but now the resummed cross section and subtracted asymptotic piece are smooth functions, which cancel well at all values of q_T (Fig. 2e-h). Hence, the distribution is physical in the whole range of q_T . Finally, at $Q^2 \gg M^2$ the small- q_T region the γ^*g fusion contribution is dominated by the $1/q_T^2$ term (Fig. 2i). Such singular terms are summed through all orders in the b -space integral corresponding to Fig. 2j and are canceled in the fusion contribution by subtracting the asymptotic piece (Fig. 2k).

To further simplify the calculation, we use the results of Refs. [12, 13], which to neglect masses of heavy quarks entering the hard scattering subprocess directly from the proton. This approximation, which differs from the complete mass-dependent cross section by higher-order terms, significantly simplifies the analysis. We therefore drop heavy-quark masses in hard scattering subdiagrams with incoming heavy quark lines. This approximation leads to massless expressions for the perturbative Sudakov factor and C -functions for quark-initiated subprocesses, and mass-dependent C -functions for gluon-initiated subprocesses.

Figure 3 demonstrates how various terms in Eq. (1) are balanced in an actual numerical calculation. This Figure shows the bottom quark production cross section vs. the polar angle θ_H of the bottom quark in the γ^*p c.m. frame for the lowest-order processes $\gamma^*b \rightarrow b$ and $\gamma^*g \rightarrow b\bar{b}$. At this order, the perturbative Sudakov factor is identically zero. Due to the heavy mass of the bottom quark, the b -space integral is practically insensitive to contributions from the nonperturbative QCD region $b \gtrsim 1 \text{ GeV}^{-1}$, so that it can be calculated without introducing a phenomenological parameterization for the Sudakov factor in the nonperturbative region. The same calculation can be done for the charm production, but in that case the nonperturbative Sudakov factor cannot be neglected. The left figure shows that near the heavy-quark mass threshold ($Q = 5 \text{ GeV}$) the b -space integral cancels well with its perturbative expansion σ_{ASY} , so that the full cross section is practically indistinguishable from the fixed-order term. On the other hand, at $Q = 50 \text{ GeV}$ (right figure) the full cross section is substantially larger than the fixed-order term at $\theta_H \lesssim 10^\circ$ and is dominated by the b -space integral. In this region, σ_{FO} is canceled well by σ_{ASY} . At $\theta_H \gtrsim 10^\circ$ the cross section agrees well with the massless result. More details can be found in our upcoming publication.

To summarize, we presented a method for the calculation of angular distributions in the heavy quark production both near the heavy-quark threshold and at large momentum transfers. Our method is realized in a massive variable flavor number factorization scheme and resums transverse momentum logarithms with the help of the b -space resummation formalism [18, 19, 20]. Our approach provides an adequate tool for detailed studies of differential distributions in a wide range of Q that will soon be available from HERA.

This work is supported by the U.S. Department of Energy, the National Science Foundation, and the Lightner-Sams Foundation.

-
- [1] J. Kroseberg (H1) (2002), hep-ex/0206042.
 - [2] W.-K. Tung, S. Kretzer, and C. Schmidt, J. Phys. **G28**, 983 (2002), hep-ph/0110247.
 - [3] M. E. Hayes and M. Kramer, J. Phys. **G25**, 1477 (1999), hep-ph/9902295.
 - [4] R. S. Thorne, J. Phys. **G25**, 1307 (1999), hep-ph/9902299.
 - [5] M. Gluck, E. Hoffmann, and E. Reya, Z. Phys. **C13**, 119 (1982).
 - [6] M. A. G. Aivazis, J. C. Collins, F. I. Olness, and W.-K. Tung, Phys. Rev. **D50**, 3102 (1994), hep-ph/9312319.
 - [7] B. A. Kniehl, M. Kramer, G. Kramer, and M. Spira, Phys. Lett. **B356**, 539 (1995), hep-ph/9505410.
 - [8] M. Buza, Y. Matiounine, J. Smith, and W. L. van Neerven, Eur. Phys. J. **C1**, 301 (1998), hep-ph/9612398.
 - [9] M. Cacciari, M. Greco, and P. Nason, JHEP **05**, 007 (1998), hep-ph/9803400.
 - [10] R. S. Thorne and R. G. Roberts, Phys. Rev. **D57**, 6871 (1998), hep-ph/9709442.
 - [11] A. Chuvakin, J. Smith, and W. L. van Neerven, Phys. Rev. **D61**, 096004 (2000), hep-ph/9910250.
 - [12] M. Kramer, F. I. Olness, and D. E. Soper, Phys. Rev. **D62**, 096007 (2000), hep-ph/0003035.
 - [13] J. C. Collins, Phys. Rev. **D58**, 094002 (1998), hep-ph/9806259.
 - [14] R. Holt et al., *The Electron Ion Collider Whitebook*, available at http://www.phenix.bnl.gov/WWW/publish/abhay/Home_of_EIC/Whitepaper/Final/.
 - [15] H. Abramowicz et al. (TESLA-N Study Group), Report DESY-01-011, Appendices, *Chapter 2: THERA: Electron proton scattering at $\sqrt{s} \approx 1 \text{ TeV}$* (2001).
 - [16] P. M. Nadolsky, N. Kidonakis, F. Olness, and C.-P. Yuan, in preparation.
 - [17] S. Kretzer, D. Mason, and F. Olness, Phys. Rev. **D65**, 074010 (2002), hep-ph/0112191.
 - [18] J. C. Collins and D. E. Soper, Nucl. Phys. **B193**, 381 (1981), erratum: *ibid.*, **B213**, 545 (1983).
 - [19] J. C. Collins and D. E. Soper, Nucl. Phys. **B197**, 446 (1982).
 - [20] J. C. Collins, D. E. Soper, and G. Sterman, Nucl. Phys. **B250**, 199 (1985).
 - [21] R. Meng, F. I. Olness, and D. E. Soper, Phys. Rev. **D54**, 1919 (1996), hep-ph/9511311.
 - [22] P. Nadolsky, D. R. Stump, and C.-P. Yuan, Phys. Rev. **D61**, 014003 (2000), erratum: *ibid.*, **D64** 059903 (2001).
 - [23] P. M. Nadolsky, D. R. Stump, and C.-P. Yuan, Phys. Rev. **D64**, 114011 (2001), hep-ph/0012261.
 - [24] H. L. Lai et al. (CTEQ), Eur. Phys. J. **C12**, 375 (2000), hep-ph/9903282.
 - [25] C. Peterson, D. Schlatter, I. Schmitt, and P. M. Zerwas, Phys. Rev. **D27**, 105 (1983).